

August 30, 2008

Comment on ‘A collider signature of the supersymmetric golden region’

Won Sang Cho, Yeong Gyun Kim and Chan Beom Park

Department of Physics, KAIST, Daejeon 305-701, Korea

Abstract

We show that Z polarization measurement is useful for distinguishing the signal decay chain of the ‘golden region’ [1] in the minimal supersymmetric standard model(MSSM) parameter space, which involves the stop decay of $\tilde{t}_2 \rightarrow \tilde{t}_1 + Z$, from other decay chains involving the neutralino decay of $\tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 + Z$ in different MSSM scenarios.

PACS numbers:

Recently, Perelstein and Spethmann investigated the ‘golden region’ in the MSSM parameter space, where the amount of fine-tuning is minimized while experimental constraints are satisfied [1]. The decay of a heavier stop into a lighter stop and a Z boson, $\tilde{t}_2 \rightarrow \tilde{t}_1 Z$, is kinematically allowed through the golden region. For instance, a benchmark point of the golden region in Ref. [1] provides stop masses, $m_{\tilde{t}_2} = 700$ GeV, $m_{\tilde{t}_1} = 400$ GeV with branching ratio $B(\tilde{t}_2 \rightarrow \tilde{t}_1 Z) = 31\%$. Then, the cascade decay in Fig.1 (a) provides a characteristic signature of the golden region scenario at the LHC. However, the interpretation of the golden region signature can be obscured by other types of cascade decays such as the one in Fig.1 (b) involving the neutralino decay $\tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 Z$, which is realized in different MSSM scenarios but produces the same final state particles as the chain in Fig.1 (a). In this note, we show that one way of distinguishing the two confusing decay chains in Fig.1, is to investigate Z boson polarization in the decay chains.




FIG. 1: (a) The decay chain of the golden region and (b) an alternative decay chain.

For the stop decay $\tilde{t}_2(p) \rightarrow \tilde{t}_1(q) + Z(k, \lambda)$, where p, q and k denote 4-momenta of the particles, and $\lambda = \pm 1, 0$ indicate the helicities of Z boson, the matrix element is given by

$$\mathcal{M} \propto (p + q)^\mu \epsilon_\mu^*(k, \lambda). \quad (1)$$

For transverse polarization vectors $\epsilon(k, \lambda)$ of Z boson, we have $p \cdot \epsilon(k, \pm 1) = q \cdot \epsilon(k, \pm 1) = 0$ in the rest frame of the decaying stop $\tilde{t}_2$, so that transverse Z polarization does not appear but only longitudinally polarized Z boson is produced from the stop decay in the rest frame. In the laboratory frame where $\tilde{t}_2$ has non-zero spatial momentum, transverse Z polarization is developed through the so-called Wigner rotation [2] even though it is absent in the stop rest

frame. However, because of large mass hierarchy between the decaying stop and Z boson for the golden region scenario, the Wigner rotation effect is so negligible that the Z boson from the stop decay in the laboratory frame is still dominantly longitudinal.

On the other hand, both of transverse and longitudinal Z polarizations are available in $\tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 + Z(\lambda)$ decay processes. In the rest frame of decaying neutralino, the relevant decay rates are given by [3]

$$\Gamma[\tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 + Z(\pm 1)] \propto m_2^2 + m_1^2 - m_Z^2 - 2m_2m_1\mathcal{A}_{\mathcal{N}}, \quad (2)$$

$$\Gamma[\tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 + Z(0)] \propto m_2^2 + m_1^2 - m_Z^2 - 2m_2m_1\mathcal{A}_{\mathcal{N}} + \frac{\lambda_Z}{m_Z^2}, \quad (3)$$

where m_1 and m_2 are the masses of $\tilde{\chi}_1^0$ and $\tilde{\chi}_2^0$ respectively, $\lambda_Z \equiv [(m_2 + m_1)^2 - m_Z^2][(m_2 - m_1)^2 - m_Z^2]$ and $\mathcal{A}_{\mathcal{N}} \equiv (|V|^2 - |A|^2)/(|V|^2 + |A|^2)$. And the vector and axial vector couplings V and A of the Z boson to the neutralino currents are expressed in terms of the neutralino mixing matrix elements [4]. Although the relative production rates of longitudinal and transverse Z polarizations in the $\tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 + Z$ decay depend on the neutralino mass spectrum, the $\tilde{\chi}_2^0\tilde{\chi}_1^0Z$ couplings and the $\tilde{\chi}_2^0$ momentum, they are expected to be similar in magnitude unless m_2 is too large compared to m_Z .




FIG. 2: The lepton angular distributions in the Z rest frame with respect to Z polarization direction for (a) the stop decay $\tilde{t}_2 \rightarrow \tilde{t}_1 + Z$ and (b) the neutralino decay $\tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 + Z$, respectively. 50 Monte Carlo events are used for the distributions.

The Z polarization can be reconstructed through lepton angular distribution in leptonic

Z decays $Z \rightarrow l^+l^-$ [3]. Fig.2 (a) shows the leptonic angular distribution of $Z \rightarrow l^+l^-$ decay in the Z rest frame with respect to the Z polarization direction, where Z bosons come from $\tilde{t}_2 \rightarrow \tilde{t}_1 + Z$ decays in the golden region scenario. On the figure, the histogram corresponds to the result of a parton-level Monte Carlo simulation with HERWIG event generator [5], while the solid line to the theoretical expectation which ignore the Wigner rotation effect. Here, 50 Monte Carlo events are used for the histogram. The number of events are expected for the benchmark point in Ref. [1] with 300 fb^{-1} luminosity if suitable event selection cuts are imposed. (See Ref.[1] for the details.) However notice that, in this work, detector response and event selection cuts are not included.

On the other hand, the corresponding lepton angular distribution for the case where Z is produced from neutralino decay of $\tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 + Z$, is shown in Fig.2 (b). Here, as a specific numerical example, we set the neutralino masses as $m_{\tilde{\chi}_2^0} = 221.6 \text{ GeV}$ and $m_{\tilde{\chi}_1^0} = 118.5 \text{ GeV}$ and assume that the neutralino mixing matrix is real. Again, on the figure, the histogram corresponds to 50 Monte Carlo events and the solid line to theoretical expectation without the Wigner rotation effect.

The two decay chains in Fig.1 exhibit very distinctive lepton angular distributions in our example so that they can be clearly distinguished experimentally, even though more careful studies including detector response, event selection cuts and the SM background would need to be performed.

In summary, we showed that Z polarization is useful to discriminate the signal decay chain of the golden region scenario which involves the stop decay $\tilde{t}_2 \rightarrow \tilde{t}_1 Z$, from other decay chains including the neutralino decay $\tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 Z$ in different MSSM scenarios.

Acknowledgments

We thank S.Y. Choi for his valuable comments on the manuscript. This work was supported by the Korea Research Foundation Grant funded by the Korean Government (MOEHRD, Basic Research Promotion Fund) (KRF-2005-210-C000006), the Center for High Energy Physics of Kyungpook National University, and the BK21 program of Ministry of

Education.

- [1] M. Perelstein and C. Spethmann, JHEP **0704** (2007) 070.
- [2] E.P. Wigner, Ann. Math. **40** (1939) 140.
- [3] S.Y. Choi and Y.G. Kim, Phys. Rev. **D 69** (2004) 015011.
- [4] S.Y. Choi, J. Kalinowski, G. Moortgat-Pick and P.M. Zerwas, Eur. Phys. J. C **22** (2001) 563
and C **23** (2002) 769; J. Kalinowski, Acta. Phys. Pol. B **34** (2003) 3441.
- [5] G. Corcella *et al.*, JHEP **01** (2001) 010, arXiv:hep-ph/0011363.